

## Universal dynamical decoupling of multiqubit states from environment

Liang Jiang<sup>1</sup> and Adilet Imambekov<sup>2</sup>

<sup>1</sup>*Institute for Quantum Information, California Institute of Technology, Pasadena, California 91125, USA*

<sup>2</sup>*Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA*

(Received 28 April 2011; published 28 December 2011)

We study the dynamical decoupling of multiqubit states from environment. For a system of  $m$  qubits, the nested Uhrig dynamical decoupling (NUDD) sequence can efficiently suppress generic decoherence induced by the system-environment interaction to order  $N$  using  $(N+1)^{2m}$  pulses. We prove that the NUDD sequence is universal, i.e., it can restore the coherence of an  $m$ -qubit quantum system independent of the details of the system-environment interaction. We also construct a general mapping between dynamical decoupling problems and discrete quantum walks in certain functional spaces.

DOI: 10.1103/PhysRevA.84.060302

PACS number(s): 03.67.Pp, 03.65.Yz, 76.60.Lz, 82.56.Jn

Dynamical decoupling (DD) is a powerful tool to protect quantum systems from decoherence induced by the inevitable system-environment interaction [1]. The idea of DD is to dynamically control the system (or environment) evolution to suppress the decoherence caused by interaction. For example, a static magnetic field with unknown magnitude  $B_z\sigma_z$  can induce dephasing of a qubit, but such dephasing can be fully eliminated by a spin flip  $\sigma_x$  (i.e., Hahn echo) at half way of the evolution [2]. In practice, however, the Hahn echo only suppresses the dephasing to  $O(T^2)$  for total evolution time  $T$ , because the magnetic field may have complicated time dependence in both magnitude and orientation due to the evolution of the environment. Furthermore, if the environment consists of quantum degrees of freedom, it can become entangled with the system via the interaction. Hence it is a challenging task to design a *universal* DD scheme that can suppress decoherence to the desired order independent of the details of the system-environment interaction.

One particularly interesting DD scheme is the concatenated DD (CDD), which has been shown to be universal for single qubits [3]. The limitation, however, is that the pulse number increases exponentially with the suppression order  $N$  [approximately  $4^N$  pulses to suppress both bit-flip and dephasing processes to  $O(T^{N+1})$ ]. It is the discovery of the universality of the Uhrig DD (UDD) sequence [4–8] that makes the universal DD practically feasible. In contrast to the CDD's demand for exponentially many pulses [3], UDD uses only  $O(N)$  spin-flip pulses to suppress the dephasing processes to  $O(T^{N+1})$  [5,6]. The discovery of the UDD sequence has inspired many experimental efforts to further improve the coherence over a wide range of quantum systems, including trapped ions [9], electron spins [10], defect centers [10,11], quantum dots [12,13], and superconducting qubits [14]. However, UDD is restricted to pure dephasing errors of a single qubit. It is desirable to have an efficient DD scheme [with  $poly(N)$  pulses] to suppress both bit-flip and dephasing processes for multiple qubits to  $O(T^{N+1})$ .

Recently, the quadratic DD (QDD) scheme has been proposed [15], which uses  $(N+1)^2$  pulses to suppress both bit-flip and dephasing errors of single qubits. As a generalization of QDD from a 1-qubit system to an  $m$ -qubit system, the nested UDD (NUDD) scheme has been proposed [16–18], which uses  $(N+1)^{2m}$  pulses to suppress decoherence from

the most general interaction between the  $m$ -qubit system and environment. Although there are numerical evidences [15] and theoretical implications [7,18,19] that QDD and NUDD might be universal, it is still an open question whether QDD and NUDD are universal or not [18,20].

In this Rapid Communication, we present a rigorous proof that the NUDD scheme with  $2m$  nesting levels and  $(N+1)^{2m}$  pulses is a universal DD scheme for an  $m$ -qubit system, which suppresses decoherence processes to  $O(T^{N+1})$  for arbitrary system-environment interaction. We achieve this by providing a mapping between NUDD and a discrete “quantum walk” in  $2m$  dimensional space. The rules that govern this quantum walk do not depend on  $2m$ , which allows an efficient proof for all nesting levels. Below we first introduce notations and explain the existing proof for UDD [6–8] using the language that can be naturally generalized for QDD and NUDD.

*UDD.* Let us first consider the UDD sequence [4] for a qubit-environment interaction

$$H(\tau) = \hat{S}_0 \otimes \hat{B}_0(\tau) + \hat{S}_1 \otimes \hat{B}_1(\tau), \quad (1)$$

where  $\hat{S}_0 = I$  and  $\hat{S}_1 = \sigma_z$ , and the time-dependent bath operators are analytic with series expansion  $\hat{B}_\alpha(t) = \sum_{p=0}^{\infty} \hat{b}_{\alpha,p} t^p$  for  $\alpha = 0, 1$ . The UDD sequence uses  $N$   $\pi$  pulses (i.e.,  $\sigma_x$  rotations) applied at times  $\tau_\lambda = T \Delta_\lambda$ , where

$$\Delta_\lambda = \sin^2 \frac{\lambda\pi}{2(N+1)} \quad (2)$$

for  $\lambda = 1, 2, \dots, N$ . (An extra  $\sigma_x$  pulse is required at  $\tau_{N+1} = T$  for  $N$  odd [15].) It is convenient to consider the toggling frame associated with the qubit. In this frame the qubit-environment Hamiltonian is modulated in time as

$$\tilde{H}(\tau) = F_0(\tau/T) S_0 \otimes \hat{B}_0(\tau) + F_1(\tau/T) S_1 \otimes \hat{B}_1(\tau), \quad (3)$$

where  $F_\alpha(t) = (-1)^{\alpha\lambda}$  for  $t \in (\Delta_\lambda, \Delta_{\lambda+1}]$ . The unitary evolution operator of the toggling frame Hamiltonian is  $\hat{U}(T) = \mathcal{T} \exp[-i \int_0^T \tilde{H}(\tau) d\tau]$ , where  $\mathcal{T}$  is the time-ordering operator.  $\hat{U}(T)$  has a Dyson expansion

$$\sum_{s=0}^{\infty} (-i)^s \sum_{\{\alpha_j, p_j\}} \hat{S}_{(\oplus \alpha_j)} \prod_{j=1}^s \hat{b}_{\alpha_j, p_j} \mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s} T^{s+p_1+\dots+p_s},$$

where  $\hat{S}_{(\oplus\alpha_j)} = \hat{S}_{\alpha_1} \cdots \hat{S}_{\alpha_s}$  and the coefficient  $\mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s}$  can be obtained by the integral [6]

$$\mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s} = \int_0^1 dt_s \cdots \int_0^{t_2} dt_1 \prod_{j=1}^s F_{\alpha_j}(t_j) t_j^{p_j}. \quad (4)$$

When  $\oplus_{j=1}^s \alpha_j = 0$ , the operator  $\hat{S}_{(\oplus\alpha_j)} = I$  is the identity operator that acts trivially on the qubit. Hence we only need to consider the terms with  $\oplus_{j=1}^s \alpha_j \neq 0$  that act nontrivially on the qubit. To show the universality of the UDD sequence, we just need to prove  $\hat{U}(T) = I \otimes \hat{U}_B(T) + O(T^{N+1})$ , which can be reduced to verifying

$$\mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s} = 0 \quad (5)$$

for  $\oplus_{j=1}^s \alpha_j \neq 0$  and  $s + \sum_{j=1}^s p_j \leq N$ . Equation (5) resembles the proof of Ref. [6] for universality of UDD. The key difference is that here additional indices  $\{\alpha_j\}$  are introduced to label different possible qubit operators ( $I$  and  $\sigma_z$ ) which will be necessary for the proof of universality of QDD and NUDD.

**QDD.** Let us now consider the QDD sequence [15] for the generic interaction between a single qubit and environment,

$$H(\tau) = \sum_{\alpha} \hat{S}_{\alpha} \otimes \hat{B}_{\alpha}(\tau), \quad (6)$$

where  $\hat{S}_{\alpha} = I, \sigma_x, \sigma_y, \sigma_z$  for binary vector labels  $\alpha = (a_2, a_1) = (0,0), (1,0), (1,1), (0,1)$ , respectively. For Pauli matrices, one can verify that  $\hat{S}_{\alpha} \hat{S}_{\alpha'} = \pm \hat{S}_{\alpha \oplus \alpha'}$ , where  $\oplus$  represents pairwise binary addition without carry [e.g.,  $(0,1) \oplus (0,1) = (0,0)$ ]. The QDD sequence consists of two nesting levels of UDD with a total of  $Q = (N+1)^2$  pulses [15]. The pulses  $\sigma_x$  and  $\sigma_z$  are associated with the first and second levels, respectively. To label these  $Q$  pulses, we introduce the vector label  $\lambda = (l_2, l_1) \in \{0, \dots, N\} \otimes \{1, \dots, N+1\}$  with  $l_2(N+1) + l_1 \in \{1, \dots, Q\}$ .<sup>1</sup> The  $\lambda$ th pulse is applied at time  $\tau_{\lambda} = T \Delta_{(l_2, l_1)}$  with

$$\Delta_{(l_2, l_1)} = \Delta_{l_2} + (\Delta_{l_2+1} - \Delta_{l_2}) \Delta_{l_1}. \quad (7)$$

The toggling frame Hamiltonian for the QDD sequence is

$$\tilde{H}(\tau) = \sum_{\alpha} F_{\alpha}(\tau/T) \hat{S}_{\alpha} \otimes \hat{B}_{\alpha}(\tau), \quad (8)$$

where  $F_{\alpha}(t) = (-1)^{a_2 l_2 + a_1 l_1}$  for  $t \in (\Delta_{(l_2, l_1)}, \Delta_{(l_2, l_1+1)})$ . One can verify that

$$F_{\alpha}(\tau/T) F_{\alpha'}(\tau/T) = F_{\alpha \oplus \alpha'}(\tau/T), \quad (9)$$

which will be useful for our proof of universality of the QDD sequence. Using the Dyson expansion of the unitary evolution operator of  $\tilde{H}(\tau)$  for QDD, we obtain that to show the suppression of both the dephasing and bit-flip errors up to  $O(T^{N+1})$  for small  $T$ , it is sufficient to prove Eq. (5) for  $\oplus_{j=1}^s \alpha_j \neq (0,0)$  and  $s + \sum_{j=1}^s p_j \leq N$ . This is very similar to UDD, the difference being that  $\alpha_j$  is now a two-component binary vector.

<sup>1</sup>We use the Greek letters  $\alpha, \beta, \lambda, \kappa$  to represent vector labels, and use the Latin letters  $a_r, b_r, l_r, k_r$  to represent the  $r$ th element of the corresponding vectors.




FIG. 1. (Color online) NUDD scheme with  $2m$  nesting levels and suppression order  $N = 2$ . (a) The timings of NUDD pulses have a self-similar structure, determined by Eqs. (2) and (10). The set of pulses associated with the  $r$ th level is  $\sigma_x^{(j)}$  for  $r = 2j - 1$ , and  $\sigma_z^{(j)}$  for  $r = 2j$ . (b) The time-dependent modulation functions  $F_{\alpha}(t)$  of the toggling frame Hamiltonian, and the corresponding pulses for the  $m$ th qubit.

**NUDD.** The NUDD sequence is a generalization of QDD from one-qubit to multiqubit systems [16–18]. For  $m$ -qubit systems, the most general system-environment interaction can be written as Eq. (6) with  $\hat{S}_{\alpha} = \sigma_{v_m}^{(m)} \otimes \sigma_{v_{m-1}}^{(m-1)} \otimes \cdots \otimes \sigma_{v_1}^{(1)}$  and  $\alpha = (a_{2m}, a_{2m-1}, \dots, a_1) \in \{0, 1\}^{\otimes 2m}$  for all generators. The Pauli operator of the  $j$ th qubit is  $\sigma_{v_j}^{(j)} = 1, \sigma_x^{(j)}, \sigma_y^{(j)}, \sigma_z^{(j)}$  for  $(a_{2j}, a_{2j-1}) = (0,0), (1,0), (1,1), (0,1)$ , respectively. The NUDD sequence consists of  $2m$  nesting levels and a total of  $Q = (N+1)^{2m}$  pulses. The decoupling pulse is  $\sigma_x^{(j)}$  for the  $(2j-1)$ th level and  $\sigma_z^{(j)}$  for the  $2j$ th level. Similar to QDD, we introduce the label  $\lambda = (l_{2m}, l_{2m-1}, \dots, l_1) \in \{0, \dots, N\}^{\otimes 2m-1} \otimes \{1, \dots, N+1\}$  with  $\sum_{r=1}^{2m} l_r (N+1)^{r-1} \in \{1, \dots, Q\}$ . The  $\lambda$ th pulse is applied at time  $\tau_{\lambda} = T \Delta_{(l_{2m}, \dots, l_1)}$ , which is defined recursively

$$\Delta_{(l_r, \dots, l_1)} = \Delta_{l_r} + (\Delta_{l_r+1} - \Delta_{l_r}) \Delta_{(l_r-1, \dots, l_1)} \quad (10)$$

for  $r = 2, \dots, 2m$ . As illustrated in Fig. 1, the timing for the pulses has a self-similar structure.<sup>2</sup> The toggling frame Hamiltonian for the NUDD sequence is the same as Eq. (8), with  $\alpha$  summed over all  $4^m$  generators. Similar to UDD and QDD, to show universality of the NUDD sequence, we just need to prove Eq. (5) for  $\oplus_{j=1}^s \alpha_j \neq \vec{0}$  and  $s + \sum_{j=1}^s p_j \leq N$ . We can view UDD and QDD as special cases of NUDD with one and two nesting levels, respectively. Since the universalities of UDD, QDD, and NUDD all rely on verifying Eq. (5), we will give a general proof of Eq. (5) in the rest of the paper.

<sup>2</sup>For  $N$  odd, pulses at different levels may coincide and form one pulse. Hence the unitary evolution of the  $\lambda$ th instantaneous pulse is  $\hat{\Omega}_{(l_{2m}, \dots, l_1 \neq N+1)} = \Omega_1$ ,  $\hat{\Omega}_{(l_{2m}, \dots, l_r \neq N, N, \dots, N, N+1)} = \Omega_r \Omega_{r-1}^N \cdots \Omega_1^N$  for  $r \geq 2$ , and  $\hat{\Omega}_{(N, \dots, N+1)} = \Omega_{2m}^N \cdots \Omega_1^N$ , where  $\Omega_{2j-1} = \sigma_x^{(j)}$  and  $\Omega_{2j} = \sigma_z^{(j)}$ .

*Universality Proof.* To prove Eq. (5), we represent each integration over  $t_j$  as a linear operator acting on functions of  $t_j$ , which generates a function of  $t_{j+1}$ . Thus, the process of multiple integrations can be thought of as a discrete quantum walk in a functional space. We choose the basis of this functional space according to the following consideration: the functional basis should be complete with respect to the operation  $\int_0^t dt' F_\alpha(t')t^p$  for all relevant orders up to  $O(T^{N+1})$ . Since  $F_\alpha(t)$  is a piecewise analytic function with at most  $Q = (N+1)^{2m}$  discontinuities, it will be convenient to use piecewise analytic functions as our functional basis. In addition, we only need to consider all polynomials up to power  $N+1$  to characterize all the effects up to  $O(T^{N+1})$ . Therefore, we choose a basis that consists of  $(N+1)Q$  functions

$$\eta_{q,\lambda}(t) = t^q \eta_\lambda(t), \quad (11)$$

with  $\eta_\lambda(t) = 1$  for  $t \in (\Delta_\lambda, \Delta_{\lambda+1}]$  and  $\eta_\lambda(t) = 0$  otherwise. Here  $q \in \{0, 1, \dots, N\}$  is the *polynomial label* and  $\lambda \in \{0, \dots, N\}^{\otimes 2m}$  is the *pulse label*.

Then, we use Eq. (9) to rewrite the integral

$$\begin{aligned} \mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s} &= \int_0^1 dt_s t_s^{p_s} \times \int_{0, [\beta_{s-1}]}^{t_s} dt_{s-1} t_{s-1}^{p_{s-1}} \\ &\dots \int_{0, [\beta_1]}^{t_2} dt_1 t_1^{p_1} \times F_{\beta_0}(t_1) \times 1, \end{aligned} \quad (12)$$

where  $\int_{0, [\beta]}^t dt' \equiv F_\beta(t) \int_0^t dt' F_\beta(t')$  and  $\beta_j = \bigoplus_{j'=1}^{s-j} \alpha_{j'}$ . For  $s + \sum_{j=1}^s p_j \leq N$ , we can compute the integral  $\mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s}$  using functional basis  $\{\eta_{q,\lambda}(t)\}$ . For each operation  $O$ , let us define the matrix form  $\mathbf{O}_{q,\lambda}^{q',\lambda'}$  as  $O \eta_{q,\lambda} = \mathbf{O}_{q,\lambda}^{q',\lambda'} \eta_{q',\lambda'}$ , where summation over repeating indices is implied. For example, the multiplication  $t \eta_{q,\lambda} = \eta_{q+1,\lambda}$  has the matrix form  $\mathbf{M}_{q,\lambda}^{q',\lambda'} = \delta_{q+1}^{q'} \delta_\lambda^{\lambda'}$ , with the Kronecker delta function  $\delta_y^x$ . The other operations are listed in Table I.

Using block diagonal properties of matrices involved,  $\mathcal{F}_{\alpha_1, \dots, \alpha_s}^{p_1, \dots, p_s}$  can be further reduced as multiplication of  $Q \times Q$  submatrices  $B_\beta$  and  $D_\beta$  [21]

$$\begin{aligned} \vec{v}_L \cdot \mathbf{M}^{p_s} \cdot \mathbf{G}_{\beta_{s-1}} \cdot \mathbf{M}^{p_{s-1}} \dots \mathbf{G}_{\beta_1} \cdot \mathbf{M}^{p_1} \cdot \mathbf{F}_{\beta_0} \vec{v}_R^T \\ = \sum_{\{i_j \geq 0\}} c_{i_1, \dots, i_s}^{\alpha_1, \alpha_2, \dots, \alpha_s} \langle u_L | D_{\beta_s}^{i_s} D_{\beta_{s-1}}^{i_{s-1}} \dots D_{\beta_1}^{i_1} B_{\beta_0} | u_R \rangle, \end{aligned}$$

where  $\sum_{j=1}^s i_j \leq s-1 + \sum_{j=1}^s p_j \leq N-1$ ,  $\beta_0 = \bigoplus_{j=1}^s \alpha_{j'}$ , and  $c_{i_1, \dots, i_s}^{\alpha_1, \alpha_2, \dots, \alpha_s}$  are possibly nonzero coefficients. The  $Q$  vectors

TABLE I. Matrix or vector forms of operations. Here  $\delta_y^x$  is the Kronecker delta function,  $B_\beta$  and  $D_\beta$  are  $Q \times Q$  matrices as defined in Eq. (13).

Operation	Matrix or vector form
$t \eta_{q,\lambda} = \eta_{q+1,\lambda}$	$\mathbf{M}_{q,\lambda}^{q',\lambda'} = \delta_{q+1}^{q'} \delta_\lambda^{\lambda'}$
$F_\beta(t) \eta_{q,\lambda} = (-1)^{\beta \lambda} \eta_{q,\lambda}$	$(\mathbf{F}_\beta)_{q,\lambda}^{q',\lambda'} = \delta_q^{q'} (B_\beta)_{\lambda,\lambda'}^{\lambda'}$
$\int_{0, [\beta]}^t dt' \eta_{q,\lambda} = (\mathbf{G}_\beta)_{q,\lambda}^{q',\lambda'} \eta_{q',\lambda'}$	$(\mathbf{G}_\beta)_{q,\lambda}^{q',\lambda'} = \delta_{q+1}^{q'} \delta_\lambda^{\lambda'} - \delta_0^{q'} (D_\beta^{q+1})_{\lambda,\lambda'}^{\lambda'}$
$\int_0^1 dt \eta_{q,\lambda} = (\vec{v}_L)_{q,\lambda}$	$(\vec{v}_L)_{q,\lambda} = \frac{\Delta_\lambda^{q+1} - \Delta_{\lambda+1}^{q+1}}{q+1}$
$1 = (\vec{v}_R^T)_{q,\lambda} \eta_{q,\lambda}$	$(\vec{v}_R^T)_{q,\lambda} = \delta_0^q$

$|u_L\rangle$  and  $|u_R\rangle$  are determined by  $\int_0^1 dt_s$  and 1 in Eq. (12), respectively, with vector elements  $|u_L\rangle_\lambda = \Delta_{\lambda+1} - \Delta_\lambda$  and  $|u_R\rangle_\lambda = 1$ . The  $Q \times Q$  matrices are

$$\begin{aligned} (B_\beta)_{\lambda,\lambda'}^{\lambda'} &= (-1)^{\beta \lambda} \delta_\lambda^{\lambda'}, \\ (D_\beta)_{\lambda,\lambda'}^{\lambda'} &= \Delta_\lambda \delta_\lambda^{\lambda'} - (\Delta_{\lambda+1} - \Delta_\lambda) \sum_{\lambda''=\lambda+1}^Q (-1)^{\beta(\lambda''-\lambda)} \delta_{\lambda''}^{\lambda'}. \end{aligned} \quad (13)$$

*Discrete quantum walk.* The last step is to verify

$$\langle u_L | D_{\beta_s}^{i_s} D_{\beta_{s-1}}^{i_{s-1}} \dots D_{\beta_1}^{i_1} | B_{\beta_0} u_R \rangle = 0 \quad (14)$$

for  $\beta_0 \neq \vec{0}$  and  $\sum_{j=1}^s i_j \leq N-1$ . The left-hand side is the amplitude of a discrete ‘‘quantum walk’’ from initial state  $|u_L\rangle$  to a target state  $|B_{\beta_0} u_R\rangle$  in the functional basis. Each multiplication of  $D_\beta$  corresponds to one step of a quantum walk. We need to show that the target state amplitude is zero when the number of steps is  $\sum_{j=1}^s i_j \leq N-1$ .

To understand the quantum walk we choose a convenient basis

$$\chi_\kappa(t) = \sum_\lambda c_{\kappa,\lambda} \eta_\lambda(t), \quad (15)$$

with the orthogonal transformation

$$c_{\kappa,\lambda} = \prod_{r=1}^{2m} \sin \left[ \frac{(2l_r + 1)k_r + (N+1)(k_{r-1} - 1) \frac{\pi}{2}}{N+1} \right], \quad (16)$$

where  $\kappa = (k_{2m}, k_{2m-1}, \dots, k_1) \in \{0, \dots, N+1\}^{\otimes 2m}$  and  $k_0 = 1$ .<sup>3</sup> In the basis  $\{|\kappa\rangle\}$  with  $|\kappa\rangle \equiv \chi_\kappa$ , the initial state is  $|u_L\rangle \propto |(1, \dots, 1)\rangle$ . Since  $\beta_0 \neq \vec{0}$  (with  $b_{r^*} = 1$  and  $b_{r < r^*} = 0$ ), the target state is  $|B_{\beta_0} u_R\rangle = \sum_{\kappa^\#} u_{\beta_0, \kappa^\#} |\kappa^\#\rangle$ , where the  $r^*$ th element of  $\kappa^\#$  is  $k_{r^*}^\# = N+1$ . After some calculation [21], it can be shown that  $D_\beta |\kappa\rangle = \sum_{k_1=0}^{k_1+1} \dots \sum_{k_{2m}=0}^{k_{2m}+1} d_{\kappa', \kappa} |\kappa'\rangle$ , where  $d_{\kappa', \kappa}$  denotes possibly nonzero coefficients. Since each step of the quantum walk only increases the index  $k_r$  by at most one unit, it requires at least  $N$  steps to walk from  $k_{r^*} = 1$  to  $k_{r^*}^\# = N+1$ . Therefore, there is zero amplitude in the target states when the number of steps  $\sum_{j=1}^s i_j \leq N-1$ . This completes the proof of Eq. (14) and implies the universality of NUDD. We emphasize that after the basis in Eq. (11) is introduced, all operations are matrix multiplications, and all our analytical statements have been explicitly checked numerically.

We can illustrate the quantum walk for special cases:

(1) For UDD, the orthogonal transformation  $c_{\kappa,\lambda} = c_{(k_1), (l_1)} = \sin[k_1 \frac{2l_1+1}{N+1} \frac{\pi}{2}]$  is simply the Fourier transformation as in Ref. [6]. The functional basis  $\{|\kappa\rangle\}$  with  $\kappa = (k_1) \in \{1, \dots, N+1\}$  forms a one-dimensional array. As illustrated in Fig. 2(a), it will take at least  $N$  steps to walk from  $\kappa = (1)$  to  $\kappa^\# = (N+1)$ . (2) For QDD, the orthogonal transformation is a little more complicated:  $c_{\kappa,\lambda} = c_{(k_2, k_1), (l_2, l_1)} = (-1)^{(k_1-1)/2} \sin[k_2 \frac{2l_2+1}{N+1} \frac{\pi}{2}] \sin[k_1 \frac{2l_1+1}{N+1} \frac{\pi}{2}]$  for odd  $k_1$  and  $(-1)^{k_1/2} \cos[k_2 \frac{2l_2+1}{N+1} \frac{\pi}{2}] \sin[k_1 \frac{2l_1+1}{N+1} \frac{\pi}{2}]$  for even  $k_1$ .

<sup>3</sup>Note that exactly  $(N+1)^{2m}$  of  $(N+2)^{2m}$  possible  $\chi_\kappa$ 's are nonzero functions, which form a new complete orthogonal basis. Other functions are simply zeros (e.g.,  $\chi_{(k_{2m}, \dots, k_2, 0)} = \chi_{(k_{2m}, \dots, k_{r+2}, N+1, 2h_r, \dots)} = \chi_{(k_{2m}, \dots, k_{r+2}, 0, 2h_r, 1, \dots)} = 0$ ), which are kept for notational convenience.




FIG. 2. (Color online) Evolution under discrete quantum walks for  $N = 4$ . The symbols represent different types of states: ‘S’ for the initial state, ‘X’ for states explored by a quantum walk, ‘#’ for unexplored target states, ‘@’ for explored target states, and ‘.’ for the remaining unexplored functional basis states. At least  $N = 4$  steps are necessary to reach the target states, which is an illustration of Eq. (14). (a) For UDD, a discrete quantum walk occurs in a one-dimensional functional basis, with initial state  $|1\rangle$ , target state  $|N + 1\rangle$ , and quantum walk matrix  $D_1$ . (b) For QDD, a discrete quantum walk occurs in a two-dimensional functional basis, with initial state  $|11\rangle$ , target states  $\{|\kappa^\#\rangle\}$ , and a quantum walk matrix  $D_{(01)}$ .

The functional basis  $\{|\kappa\rangle\}$  with  $\kappa = (k_2, k_1)$  forms a two-dimensional array. As illustrated in Fig. 2(b), it will also take at least  $N$  steps to walk from  $\kappa = (1, 1)$  to  $\kappa^\# = (N + 1, k_1)$  or  $(k_2, N + 1)$ . (3) For NUDD, the functional basis forms a  $2m$ -dimensional array. Similar to UDD and QDD, it will take at least  $N$  steps to walk from  $\kappa = (1, \dots, 1)$  to  $\kappa^\#$  with  $k_r^\# = N + 1$ .

When the suppression order is  $N_r$  for the  $r$ th nesting level of NUDD, the overall suppression of decoherence is  $O(T^{N^*+1})$  limited by the lowest suppression order  $N^* = \min[N_1, \dots, N_{2m}]$ .

*Summary and outlook.* We have proved the universality of the NUDD sequence, which can restore an unknown initial state of an  $m$ -qubit system to  $O(T^{N+1})$  using  $2m$  nesting levels and  $(N + 1)^{2m}$  pulses, independent of the details of the system-environment interaction. The NUDD sequence is

experimentally feasible, because it requires only  $poly(N)$  pulses acting on individual qubits. Our work illustrates a general connection between DD problems and discrete quantum walks. The techniques developed can be used to address a variety of questions, such as investigation of environment correlations using DD, schemes of efficient DD for particular system-environment interactions, and the combination of multiqubit DD schemes with quantum error correcting codes [22] and quantum algorithms.

*Note added.* Recently, a preprint of closely related work by Kuo and Lidar [23] became available, with a different proof of the universality of QDD.

*Acknowledgements.* We thank John Preskill for helpful discussions. This work was supported by the Sherman Fairchild Foundation, A.P. Sloan Foundation under Grant No. BR-5123, and NSF Career Award DMR-1049082.

- 
- [1] L. Viola, E. Knill, and S. Lloyd, *Phys. Rev. Lett.* **82**, 2417 (1999).
  - [2] E. L. Hahn, *Phys. Rev.* **80**, 580 (1950).
  - [3] K. Khodjasteh and D. A. Lidar, *Phys. Rev. Lett.* **95**, 180501 (2005).
  - [4] G. S. Uhrig, *Phys. Rev. Lett.* **98**, 100504 (2007).
  - [5] B. Lee, W. M. Witzel, and S. Das Sarma, *Phys. Rev. Lett.* **100**, 160505 (2008).
  - [6] W. Yang and R.-B. Liu, *Phys. Rev. Lett.* **101**, 180403 (2008).
  - [7] S. Pasini and G. S. Uhrig, *J. Phys. A* **43**, 132001 (2010).
  - [8] G. S. Uhrig and D. A. Lidar, *Phys. Rev. A* **82**, 012301 (2010).
  - [9] M. J. Biercuk, H. Uys, A. P. VanDevender, N. Shiga, W. M. Itano, and J. J. Bollinger, *Nature (London)* **458**, 996 (2009).
  - [10] G. de Lange, Z. H. Wang, D. Riste, V. V. Dobrovitski, and R. Hanson, *Science* **330**, 60 (2010).
  - [11] C. A. Ryan, J. S. Hodges, and D. G. Cory, *Phys. Rev. Lett.* **105**, 200402 (2010).
  - [12] H. Bluhm, S. Foletti, I. Neder, M. Rudner, D. Mahalu, V. Umansky, and A. Yacoby, *Nat. Phys.* **7**, 109 (2011).
  - [13] C. Barthel, J. Medford, C. M. Marcus, M. P. Hanson, and A. C. Gossard, *Phys. Rev. Lett.* **105**, 266808 (2010).
  - [14] J. Bylander, S. Gustavsson, F. Yan, F. Yoshihara, K. Harrabi, G. Fitch, D. G. Cory, Y. Nakamura, J.-S. Tsai, and W. D. Oliver, *Nat. Phys.* **7**, 565 (2011).
  - [15] J. R. West, B. H. Fong, and D. A. Lidar, *Phys. Rev. Lett.* **104**, 130501 (2010).
  - [16] M. Mukhtar, T. B. Saw, W. T. Soh, and J. Gong, *Phys. Rev. A* **81**, 012331 (2010).
  - [17] M. Mukhtar, W. T. Soh, T. B. Saw, and J. Gong, *Phys. Rev. A* **82**, 052338 (2010).
  - [18] Z.-Y. Wang and R.-B. Liu, *Phys. Rev. A* **83**, 022306 (2011).
  - [19] Z.-Y. Wang and R.-B. Liu, *Phys. Rev. A* **83**, 062313 (2011).
  - [20] W. Yang, Z.-Y. Wang, and R.-B. Liu, *Front. Phys. China* **6**, 2 (2011).
  - [21] L. Jiang and A. Imambekov (unpublished).
  - [22] H. K. Ng, D. A. Lidar, and J. Preskill, *Phys. Rev. A* **84**, 012305 (2011).
  - [23] W.-J. Kuo and D. A. Lidar, *Phys. Rev. A* **84**, 042329 (2011).